

NMT- Neuromuscular Transmission Publications Reference List

January 2017

Contents

Articles on research conducted with GE Healthcare NMT modules	2
Articles on research conducted with other or unknown equipment.....	4
General and editorial articles & publications	6

Articles on research conducted with GE Healthcare NMT modules

- Kaufhold N., et al. Sugammadex and neostigmine dose-finding study for reversal of residual neuromuscular block at a train-of-four ratio of 0.2 (SUNDRO20). *British Journal of Anaesthesia*, **116(2)**, 233-240 (2016).
- Salminen J., et al. Comparison of train-of-four ratios measured with Datex-Ohmeda's M-NMT MechanoSensor™ and M-NMT ElectroSensor™. *Journal of Clinical Monitoring and Computing*, **30(3)**, 295-300 (2016).
- Ghaffar E.A., et al. Kinemyography (KMG) Versus Electromyography (EMG) Neuromuscular Monitoring in Pediatric Patients Receiving Rocuronium during General Anesthesia *Med. J. Cairo Univ.*, **83(1)**, 121-128 (2015).
- Bezen E.F., et al. Comparison of acceleromyography and kinemyography monitoring methods during recovery period of neuromuscular block. *European Journal of Anaesthesiology*, **3AP3-9, 31**, 38 (2014).
- Stewart P. A., et al. Comparison of electromyography and kinemyography during recovery from non-depolarising neuromuscular blockade. *Anaesthesia and Intensive Care*, **42(3)**, 378 (2014).
- Todd S.S., et al. The implementation of Quantitative Electromyographic Neuromuscular Monitoring in an Academic Anesthesia Department. *Anesthesia & Analgesia*, **119(2)**, 324-331 (2014).
- Kim W.H., et al. Comparison of motor-evoked potentials monitoring in response to transcranial electrical stimulation in subjects undergoing neurosurgery with partial vs no neuromuscular block. *British Journal of Anaesthesia*, **110(4)**, 567-576 (2013).
- Liang S. S., et al. An ipsilateral comparison of acceleromyography and electromyography during recovery from nondepolarizing neuromuscular block under general anesthesia in humans. *Anesthesia & Analgesia*, **117(2)**, 373-379 (2013).
- Nitahara K., et al. Recovery of train-of-four ratio to 0.70 and 0.90 is delayed in type 2 diabetes with vecuronium-induced neuromuscular block. *European Journal of Anaesthesiology (EJA)*, **30(2)**, 80-84 (2013).
- Sugi Y., et al. Acceleromyography at the Flexor Hallucis Brevis Muscle Underestimates Residual Neuromuscular Blockade. *The Open Anesthesiology Journal*, **7**, 26-29 (2013).
- Ghodraty M. R., et al. The effects of magnesium sulfate on neuromuscular blockade by cisatracurium during induction of anesthesia. *Journal of Anesthesia*, **26(6)**, 858-863 (2012).
- Hans G.A., et al. Intravenous magnesium re-establishes neuromuscular block after spontaneous recovery from an intubating dose of rocuronium: a randomised controlled trial. *European Journal of Anaesthesiology (EJA)*, **29(2)**, 95-99 (2012).
- S. Phillips, P. A. Stewart, N. Freeland, G. Heller. Comparison of evoked electromyography in three muscles of the hand during recovery from non-depolarising neuromuscular blockade. *Anaesth Intensive Care*; **40**: 690-696 (2012)
- Paul A Stewart, Min Li Huang, Susan Qiushuang Li, Ayse Bilgin, Stephanie Phillips. Critical respiratory events and residual neuromuscular blockade in the postoperative recovery unit in 2012. *ANZCA ASM 2012*
- Deckert D., et al. Unexpected failure of rocuronium-mediated neuromuscular blockade. *Anaesthesia*, **66(1)**, 58-59 (2011).
- Sugi Y., et al. Comparison of the Neuromuscular Profile of the First Dorsal Interosseous Muscle and the Flexor Hallucis Brevis Muscle as Measured by Electromyography. *ISRN Anesthesiology* (2011).
- Brull, S.J. and Murphy G.S., Residual neuromuscular block: lessons unlearned. Part II: methods to reduce the risk of residual weakness. *Anesthesia & Analgesia* **111(1)** 129-140 (2010).
- Murphy G.S. and Brull S.J., Residual neuromuscular block: lessons unlearned. Part I: definitions, incidence, and adverse physiologic effects of residual neuromuscular block. *Anesthesia & Analgesia*, **111(1)**, 120-128, (2010).
- Hans G.A., et al. Effect of an intravenous infusion of lidocaine on cisatracurium induced neuromuscular block duration: a randomized controlled trial. *Acta Anaesthesiologica Scandinavica*, **54(10)**, 1192-1196 (2010).
- Saitoh Y., et al. Monitoring of vecuronium-induced neuromuscular block at the sternocleidomastoid muscle in anesthetized patients. *Journal of Anesthesia*, **24(6)**, 838-844 (2010).
- Adamus M., et al. Influence of gender on the course of neuromuscular block following a single bolus dose of cisatracurium or rocuronium. *European Journal of Anaesthesiology*, **25(7)**, 589-595 (2008).

- Dahaba A. A., et al. Comparison of a new neuromuscular transmission monitor compressomyograph with mechanomyograph. *British Journal of Anaesthesia*, **100(3)**, 344-350 (2008).
- Gilhuly T.J., et al. Improved neuromuscular blockade using a novel neuromuscular blockade advisory system: a randomized, controlled, clinical trial. *Anesthesia & Analgesia*, **107(5)**, 1609-1617 (2008).
- Motamed C. and Bourgain J-L., Computerized recording of neuromuscular monitoring and the risk of residual paralysis at the time of extubation. *Journal of Clinical Monitoring and Computing*, **22(4)**, 315-318 (2008).
- Saitoh Y., et al. Recovery of neuromuscular blockade caused by vecuronium is delayed in patients with hypertriglyceridemia. *Journal of Anesthesia*, **22(1)**, 89-92 (2008).
- Saitoh Y., et al. Monitoring of vecuronium-induced neuromuscular blockade during one-lung ventilation. *Journal of Anesthesia*, **22(4)**, 378-384 (2008).
- Adamus M., et al. Influence of Gender on the Onset and Duration of Rocuronium-induced Neuromuscular Block. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub*, **151(2)**, 301-305 (2007).
- Adamus M. and Belohlavek R., Fuzzy control of neuromuscular block during general anesthesia-system design, development and implementation. *International Journal of General Systems*, **36(6)**, 733-743 (2007).
- Hänzi P., et al. Clinical validation of electromyography and acceleromyography as sensors for muscle relaxation *European Journal of Anaesthesiology*, **24**, 882-888 (2007).
- Schumacher P. M., et al. Model-based control of neuromuscular block using mivacurium: design and clinical verification. *European Journal of Anaesthesiology*, **23(8)**, 691-699 (2006).
- Trager G., et al. Comparison of phonomyography, kinemyography and mechanomyography for neuromuscular monitoring. *Canadian Journal of Anesthesia*, **53(2)**, 130-135 (2006).
- Adamus M., et al. Predictability of rocuronium-induced neuromuscular block: A-456. *European Journal of Anaesthesiology (EJA)*, **22**, 120 (2005).
- Kopman A. F., et al. Acceleromyography vs. electromyography: an ipsilateral comparison of the indirectly evoked neuromuscular response to train-of-four stimulation. *Acta Anaesthesiologica Scandinavica*, **49(3)**, 316-322 (2005).
- Richard A., et al. Cisatracurium-induced neuromuscular blockade is affected by chronic phenytoin or carbamazepine treatment in neurosurgical patients. *Anesthesia & Analgesia*, **100(2)**, 538-544 (2005).
- Vasella F.C., et al. Antagonism of neuromuscular blockade but not muscle relaxation affects depth of anaesthesia. *British Journal of Anaesthesia*, **94(6)**, 742-747 (2005).
- Mendonça T., et al. Hipocrates: a robust system for the control of neuromuscular blockade. *Journal of Clinical Monitoring and Computing*, **18(4)**, 265-273 (2004).
- Motamed C., et al. Comparison between the Datex-Ohmeda M-NMT® module and a force-displacement transducer for monitoring neuromuscular blockade. *European Journal of Anaesthesiology*, **20(6)**, 467-469 (2003).
- Saitoh Y., et al. Monitoring of neuromuscular block after administration of vecuronium in patients with diabetes mellitus. *British Journal of Anaesthesia*, **90(4)**, 480-486 (2003).
- Dahaba A. A., et al. The neuromuscular transmission module versus the relaxometer mechanomyograph for neuromuscular block monitoring. *Anesthesia & Analgesia*, **94(3)**, 591-596 (2002).
- Dahaba A.A., et al. AS/3® neuromuscular transmission module (M-NMT) vs. mechanomyography for neuromuscular block monitoring. *European Journal of Anaesthesiology*, **19(3)**, 210-211 (2002).
- Goldhill D.R., et al. Neostigmine after spontaneous recovery from neuromuscular blockade. *Anaesthesia*, **44(4)**, 293-299 (1989).

Articles on research conducted with other or unknown equipment

- Grosse-Sundrup M., et al. Intermediate acting non-depolarizing neuromuscular blocking agents and risk of postoperative respiratory complications: prospective propensity score matched cohort study. *British Medical Journal*, **345**, e6329 (2012).
- Murphy GS, Szokol JW, Marymont JH, Greenberg SB, Avram MJ, Vender JS. Residual neuromuscular blockade and critical respiratory events in the postanesthesia care unit. *Anesthesia & Analgesia*, **107(1)**:130-7 (2008)
- Naguib M., et al. Neuromuscular monitoring and postoperative residual curarisation: a meta-analysis. *British Journal of Anaesthesia*, **98(3)**, 302-316 (2007).
- Suzuki T., et al. Epidurally administered mepivacaine delays recovery of train-of-four ratio from vecuronium-induced neuromuscular block. *British Journal of Anaesthesia*, **99(5)**, 721-725 (2007).
- Baillard C., et al. Postoperative residual neuromuscular block: a survey of management. *British Journal of Anaesthesia*, **95(5)**, 622-626 (2005).
- Hattori H., et al. Gabexate mesilate hastens recovery from vecuronium-induced neuromuscular blockade. *European Journal of Anaesthesiology*, **22(1)**, 20-24 (2005).
- Berg, H, et al. Residual neuromuscular block is a risk factor for postoperative pulmonary complications, *Acta Anaesthesiologica Scandinavica*. **41**, 1095-1103 (1997).
- Baurain M.J., et al. Is recovery of neuromuscular transmission complete after the use of neostigmine to antagonize block produced by rocuronium, vecuronium, atracurium and pancuronium? *British Journal of Anaesthesia*, **77(4)**, 496-499 (1996).
- Kirkegaard-Nielsen H., et al. Response to double-burst appears before response to train-of-four stimulation during recovery from non-depolarizing neuromuscular blockade. *Acta Anaesthesiologica Scandinavica*, **40(6)**, 719-723 (1996).
- Debaene B., et al. Monitoring the onset of neuromuscular block at the orbicularis oculi can predict good intubating conditions during atracurium-induced neuromuscular block. *Anesthesia & Analgesia*, **80(2)**, 360-363 (1995).
- McCoy É.P., et al. Nondepolarizing neuromuscular blocking drugs and train-of-four fade. *Canadian Journal of Anaesthesia*, **42(3)**, 213-216 (1995).
- Mortensen C. R. et al. Perioperative monitoring of neuromuscular transmission using acceleromyography prevents residual neuromuscular block following pancuronium. *Acta Anaesthesiologica Scandinavica*, **39(6)**, 797-801 (1995).
- Helbo-Hansen HS, Bang U, Nielsen HK, Skovgaard LT. The accuracy of train-of-four monitoring at varying stimulating currents. *Anesthesiology*. 1992;76 (2):199-203
- Baxter M.R.N., et al. Postoperative neuromuscular function in pediatric day-care patients. *Anesthesia & Analgesia*, **72(4)**, 504-508 (1991).
- Brull, S. et al. Effect of tetanus on subsequent neuromuscular monitoring in patients receiving vecuronium. *Anesthesiology* **74**, 64-70 (1991).
- Donati F., et al. Vecuronium neuromuscular blockade at the adductor muscles of the larynx and adductor pollicis. *Anesthesiology*, **74(5)**, 833-837 (1991).
- Beemer, G. et al. Accurate monitoring of neuromuscular blockade using a peripheral nerve stimulator – a review. *Anaesth Intens Care* **18**, 490-496 (1990).
- Donati F., et al. Vecuronium neuromuscular blockade at the diaphragm, the orbicularis oculi, and adductor pollicis muscles. *Anesthesiology*, **73(5)**, 870-875 (1990).
- Toft P., et al. Effect of epidurally administered bupivacaine on atracurium-induced neuromuscular blockade. *Acta Anaesthesiologica Scandinavica*, **34(8)**, 649-652 (1990).
- Engbaek J., et al. Double burst stimulation (DBS): a new pattern of nerve stimulation to identify residual neuromuscular block. *British Journal of Anaesthesia*, **62(3)**, 274-278 (1989).

- Gwinnutt C. L. and Meakin G., Use of the post-tetanic count to monitor recovery from intense neuromuscular blockade in children. *British Journal of Anaesthesia*, **61(5)**, 547-550 (1988).
- Ridley S.A. and Hatch D.J., Post-tetanic count and profound neuromuscular blockade with atracurium infusion in paediatric patients. *British Journal of Anaesthesia*, **60(1)**, 31-35 (1988).
- Bonsu A.K., et al. Relationship of post-tetanic count and train-of-four response during intense neuromuscular blockade caused by atracurium. *British Journal of Anaesthesia*, **59(9)**, 1089-1092 (1987).
- Chauvin M., et al. The neuromuscular blocking effect of vecuronium on the human diaphragm. *Anesthesia & Analgesia* **66(2)**, 117-122, (1987).
- Curran M.J., et al. Onset and recovery of Atracurium and Suxsmethonium-induced neuromuscular blockade with simultaneous Train-Of-Four and single twitch stimulation. *British Journal of Anaesthesia*, **59(8)**, 989-994 (1987).
- Fernando P.U.E., et al. Relationship between posttetanic count and response to carinal stimulation during vecuronium-induced neuromuscular blockade. *Acta Anaesthesiologica Scandinavica*, **31(7)**, 593-596 (1987).
- Ornstein E., et al. The effect of phenytoin on the magnitude and duration of neuromuscular block following atracurium or vecuronium. *Anesthesiology*, **67(2)**, 191-196 (1987).
- Beemer, G. et al. Postoperative neuromuscular function. *Anaesthesia & Intensive Care* **14**, 41-45 (1986).
- Pearce A.C., et al. Factors affecting train-of-four fade. *British Journal of Anaesthesia*, **57(6)**, 602-606 (1985).
- Lee C., and DeSilva A.J.C. Acute and subchronic neuromuscular blocking characteristics of streptomycin: A comparison with neomycin. *British Journal of Anaesthesia*, **51(5)**, 431-434 (1979).
- Lee C., et al. Neuromuscular sensitivity to tubocurarine A comparison of 10 parameters. *British Journal of Anaesthesia*, **48(11)**, 1045-1051 (1976).
- Ali H.H., et al. The effect of tubocurarine on indirectly elicited train-of-four muscle response and respiratory measurements in humans. *British Journal of Anaesthesia*, **47(5)**, 570-574 (1975).
- Lee C. Train-of-4 quantitation of competitive neuromuscular block. *Anesthesia & Analgesia*, **54(5)**, 649-653 (1975).
- Ali, H. *et al.* Evaluation of recovery from nondepolarizing neuromuscular block, using a digital neuromuscular transmission analyzer: preliminary report. *Anesth Analg* **52**, 740-745 (1973).
- Ali H.H., *et al.* Stimulus frequency in the detection of neuromuscular block in humans. *British Journal of Anaesthesia*, **42(11)**, 967-978 (1970).
- Ghoneim M. M. and Long L.P. The interaction between magnesium and other neuromuscular blocking agents. *The Journal of the American Society of Anesthesiologists*, **32(1)**, 23-26 (1970).
- Katz R.L., Neuromuscular effects of d-tubocurarine, edrophonium and neostigmine in man. *The Journal of the American Society of Anesthesiologists*, **28(2)**, 327-336 (1967).
- Deacock A.R. de C. and Hargrove R.L., The influence of certain ganglionic blocking agents on neuromuscular transmission. *British Journal of Anaesthesia*, **34(6)**, 357-362 (1962).

General and editorial articles & publications

- M. Naguib, S. J. Brull and K. B. Johnson, Conceptual and technical insights into the basis of neuromuscular monitoring. *Anaesthesia*, **72** (Suppl. 1), 16–37 (2017)
- Renew J.R. and Brull S.J., The Effect of Quantitative Neuromuscular Monitoring on the Incidence of Residual Neuromuscular Blockade and Clinical Outcomes. *Current Anesthesiology Reports*, **6(2)**, 170-177 (2016).
- Stoelting, R. K. Monitoring of neuromuscular blockade: what would you expect if you were the patient. *APSF Newsletter*, **30**, 45-7 (2016)
- Checketts M.R., *et al.*, Recommendations for standards of monitoring during anaesthesia and recovery 2015: Association of Anaesthetists of Great Britain and Ireland
- Lien, C. A., & Kopman, A. F. Current recommendations for monitoring depth of neuromuscular blockade. *Current Opinion in Anesthesiology*, **27(6)**, 616-622 (2014)
- Hemmerling T., Neuromuscular Monitoring Part 1: A Review and Update *Anesthesiology Rounds*, **3(3)**, (2013).
- Loughnan T.E. and Loughnan A.J. , Overview of the introduction of neuromuscular monitoring to clinical anaesthesia. *Anaesthesia and Intensive Care* (**41**)**1**, 19 (2013).
- Dubois P. E., *et al.* Datex-Ohmeda NeuroMuscular Transmission electromyography module artefacts in clinical practice: case report and retrospective chart review. *European Journal of Anaesthesiology (EJA)*, **29(5)**, 249-251 (2012).
- Kopman A.F. Monitoring Neuromuscular Function: Facts and Fallacies. *ASA Refresher Courses in Anesthesiology*, **39(1)**, 72-79 (2011).
- Kopman A.F., Perioperative monitoring of neuromuscular function. *Monitoring in Anesthesia and Perioperative Care*, **22**, 261-273 (2011).
- Fuchs-Buder T. Principles of neuromuscular monitoring. *Neuromuscular monitoring in clinical practice and research, Springer Berlin, Heidelberg*, 24-72 (2010).
- Fagerlund M.J. and Eriksson L.I., Current concepts in neuromuscular transmission. *British Journal of Anaesthesia*, **103(1)**, 108-114 (2009).
- Martyn J.A.J., M. *et al.* Basic principles of neuromuscular transmission. *Anaesthesia* **64(1)**, 1-9 (2009).
- Srivastava A. and Hunter J.M., Reversal of neuromuscular block. *British Journal of Anaesthesia*, **103(1)**, 115-129 (2009).
- Fuchs-Buder T., *et al.* Good clinical research practice in pharmacodynamic studies of neuromuscular blocking agents II: the Stockholm revision. *Acta Anaesthesiologica Scandinavica*, **51(7)**, 789-808 (2007).
- Hemmerling, T.M. & Le, N. Brief review: Neuromuscular monitoring: an update for the clinician, *Can J Anesth* (2007) 54: 58. doi:10.1007/BF03021901
- Kopman, A. F. The Datex-Ohmeda M-NMT module: a potentially confusing user interface. *The Journal of the American Society of Anesthesiologists*, **104(5)**, 1109-1110 (2006).
- McGrath C. D. and Hunter J.M., Monitoring of neuromuscular block. *Continuing Education in Anaesthesia, Critical Care & Pain*, **6(1)**, 7-12 (2006).
- Trager, G., *et al.* Comparison of phonomyography, kinemyography and mechanomyography for neuromuscular monitoring. *Canadian Journal of Anesthesia* **53(2)**, 130-135 (2006).
- Kopman A. F., Is the Datex-Ohmeda® M-NMT module a research quality instrument? *European Journal of Anaesthesiology*, **21(10)**, 833-834 (2004).
- Nakamura K., *et al.* Erythemas caused by electrodes while monitoring neuromuscular blockade: three cases. *Journal of Anesthesia*, **18(4)**, 296-299 (2004).

- El-Orbany M.I., et al. The relationship of posttetanic count and train-of-four responses during recovery from intense cisatracurium-induced neuromuscular blockade. *Anesthesia & Analgesia*, **97(1)**, 80-84 (2003).
- Dahaba A. A., The M-NMT Mechanosensor Cannot be Considered as a Reliable Clinical Neuromuscular Monitor in Daily Anesthesia Practice. *Anesthesia & Analgesia*, **95(6)**, 1827 (2002).
- Hemmerling T. M. and Donati F., The M-NMT mechanosensor cannot be considered as a reliable clinical neuromuscular monitor in daily anesthesia practice. *Anesthesia & Analgesia*, **95(6)**, 1826-1827 (2002).
- Viby-Mogensen J., Neuromuscular monitoring. *Current Opinion in Anesthesiology* **14(6)**, 655-659 (2001).
- Fisher D.M., Neuromuscular blocking agents in paediatric anaesthesia. *British Journal of Anaesthesia*, **83(1)**, 58-64 (1999).
- Eriksson LI, Sundman E. et al. Functional assessment of the pharynx at rest and during swallowing in partially paralyzed humans: simultaneous videomanometry and mechanomyography of awake human volunteers. *Anesthesiology*, **87(5)**: 1035-104 (1997)
- Rowaan C.J., et al. The Relaxometer: a complete and comprehensive computer-controlled neuromuscular transmission measurement system developed for clinical research on muscle relaxants. *Journal of Clinical Monitoring*, **9(1)**, 38-44 (1993).
- Westenskow D.R. and Joseph A., Method and apparatus for monitoring neuromuscular blockage. *U.S. Patent*, **5(131)**, 401 (1992).
- Jensen E., et al. The Accelograph®: a new neuromuscular transmission monitor. *Acta Anaesthesiologica Scandinavica* **32(1)**, 49-52 (1988).
- Jones R.M., Neuromuscular transmission and its blockade. *Anaesthesia*, **40(10)**, 964-976 (1985).
- Viby-Mogensen J. Clinical assessment of neuromuscular transmission. *British Journal of Anaesthesia*, **54(2)**, 209-223 (1982).
- Ali H.H., et al. Twitch, tetanus and train-of-four as indices of recovery from nondepolarizing neuromuscular blockade. *Anesthesiology* **54**, 294-297 (1981).
- Standaert F.G. and Dretchen K.L., Cyclic nucleotides in neuromuscular transmission. *Anesthesia & Analgesia*, **60(2)**, 91-99 (1981).
- Bowman W.C., Prejunctional and Postjunctional Cholinceptors at the Neuromuscular Junction. *Anesthesia & Analgesia*, **59(12)**, 935-943 (1980).
- Argov, Z., et al. Disorders of neuromuscular transmission caused by drugs. *N Engl J Med* **301**, 409-413 (1979).
- Brand J.B., et al. Spontaneous recovery from nondepolarizing neuromuscular blockade: correlation between clinical and evoked responses. *Anesthesia & Analgesia*, **56(1)**, 55-58 (1977).
- Ali H.H., et al. Monitoring of neuromuscular function. *Anesthesiology* **45**, 216-249 (1976).
- Lee C.-M., Train-of-4 quantitation of competitive neuromuscular block. *Anesthesia & Analgesia*, **54(5)**, 649-653 (1975).



Imagination at work

Product may not be available in all countries and regions. Full product technical specification is available upon request. Contact a GE Healthcare Representative for more information. Please visit www.gehealthcare.com/promotional-locations.

Data subject to change.

© 2017 General Electric Company – All rights reserved.

GE, the GE monogram and imagination at work are trademarks of General Electric Company.

GE Healthcare reserves the right to make changes in specifications and features shown herein, or discontinue the product described at any time without notice or obligation. Contact your GE Healthcare representative for the most current information.

GE Healthcare, a division of General Electric Company.

Reproduction in any form is forbidden without prior written permission from GE. Nothing in this material should be used to diagnose or treat any disease or condition. Readers must consult a healthcare professional.

JB45670XX 1/17